



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Similarly, $JI'' \perp CA$ and $JI''' \perp AB$.

(4) Arc $P''B = \text{arc } P'C$. Hence, $\angle P''PB = \angle P'PC$; hence $\angle BPR = \angle RPC$. Hence, $BR : RC = BP : PC$. Similarly, $CR' : R'A = CP' : P'A$, and $AR'' : R''B = AP'' : P''B$. Hence,

$$\frac{BR \cdot CR' \cdot AR''}{RC \cdot R'A \cdot R''B} = \frac{BP \cdot CP' \cdot AP''}{PC \cdot P'A \cdot P''B}.$$

But $BP = AP'$, being opposite sides of a rectangle. Similarly, $CP' = BP''$ and $AP'' = CP$. Hence,

$$\frac{BR \cdot CR' \cdot AR''}{RC \cdot R'A \cdot R''B} = 1,$$

numerically. Hence, AR, BR', CR'' are concurrent.

Also solved by A. M. HARDING.

CALCULUS.

373. Proposed by C. N. SCHMALL, New York City.

In the *Encyclopaedia Britannica* article on "Capillary Action" (Vol. 5, p. 268, 11th ed.) it is shown that $1/R_1 + 1/R_2 = p/T$, in the case of a soap bubble, where R_1, R_2 are the *principal radii of curvature* at any point of the bubble; p , the difference of air-pressure; T , the energy per unit area of the film. Employing the principle that the soap bubble tends to assume a form such that the area of its surface is a *minimum* for a *given volume* of air, show by the calculus of variations that $1/R_1 + 1/R_2 = k$, a constant.

SOLUTION BY THE PROPOSER.

We have here to determine the solid which, with a given volume (capacity), contains the least surface. Hence, we have to make the surface integral

$$U = \iint \sqrt{1 + p^2 + q^2} \, dxdy \quad (1)$$

a minimum, subject to the condition that the volume integral

$$I = \iint z \, dxdy \quad (2)$$

is constant.

It should be remembered that in this discussion $p = \partial z / \partial x$, $q = \partial z / \partial y$, $r = \partial^2 z / \partial x^2$, $s = \partial^2 z / \partial x \partial y$, and $t = \partial^2 z / \partial y^2$.

Let k be a constant. Then it is evident that, when the surface (1) is a minimum, the binomial

$$\begin{aligned} \iint \sqrt{1 + p^2 + q^2} \, dxdy + \iint kz \, dxdy &\equiv \iint (\sqrt{1 + p^2 + q^2} + kz) \, dxdy \\ &\equiv \iint V \, dxdy \end{aligned} \quad (3)$$

will also be a minimum.

Now, taking x and y as the independent variables, the condition for a mini-

num is, by the calculus of variations,

$$\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} = L, \quad (4)$$

(TODHUNTER'S *Int. Cal.*, p. 374, 5th ed.)

where

$$M = \frac{\partial V}{\partial p} = \frac{p}{\sqrt{1+p^2+q^2}},$$

$$N = \frac{\partial V}{\partial q} = \frac{q}{\sqrt{1+p^2+q^2}},$$

$$L = \frac{\partial V}{\partial z} = k.$$

(TODHUNTER'S, *Int. Cal.*, p. 372.)

Hence, equation (4) becomes

$$\frac{\partial}{\partial x} \left(\frac{p}{\sqrt{1+p^2+q^2}} \right) + \frac{\partial}{\partial y} \left(\frac{q}{\sqrt{1+p^2+q^2}} \right) = k \quad (5)$$

or,

$$\frac{r(1+p^2+q^2) - (pr+qs)p + t(1+p^2+q^2) - (ps+qt)q}{(1+p^2+q^2)^{\frac{3}{2}}} = k,$$

or,

$$\frac{(1+q^2)r - 2pqs + (1+p^2)t}{(1+p^2+q^2)^{\frac{3}{2}}} = k, \quad (6)$$

which is the partial differential equation of the required minimal surface, the integral of which will represent the surface itself.

Again, R_1 and R_2 are known as the *principal radii of curvature* at any point of the surface. The equation giving these is

$$(rt - s^2)R^2 - \sqrt{1+p^2+q^2}[(1+p^2)t - 2pqs + (1+q^2)r]R + (1+p^2+q^2)^2 = 0. \quad (7)$$

(GOURSAT-HEDRICK'S *Math. Anal.*, Vol. 1, p. 504, Eq. 13.)

If R_1, R_2 , be the roots of this quadratic in R , we have

$$R_1 + R_2 = \frac{\sqrt{1+p^2+q^2}[(1+p^2)t - 2pqs + (1+q^2)r]}{rt - s^2}, \quad (8)$$

$$R_1 R_2 = \frac{(1+p^2+q^2)^2}{rt - s^2}. \quad (9)$$

Dividing (8) by (9), we obtain

$$\frac{1}{R_1} + \frac{1}{R_2} = \frac{(1+p^2)t - 2pqs + (1+q^2)r}{(1+p^2+q^2)^{\frac{3}{2}}}. \quad (10)$$

(See EISENHART'S *Diff. Geom.*, p. 126, ex. 3.)

Comparing equations (6) and (10) we have the required result,

$$\frac{1}{R_1} + \frac{1}{R_2} = k.$$

Note 1.—The sphere,

$$x^2 + y^2 + z^2 = a^2 \quad (11)$$

and the cylinder,

$$y^2 + z^2 = b^2, \quad (12)$$

are examples of minimal surfaces satisfying Eq. (6).

Thus, in (11),

$$p = -x/z, \quad q = -y/z, \quad \sqrt{1+p^2+q^2} = a/z,$$

$$\therefore M = -x/a, \quad N = -y/a,$$

and (11) becomes $k + 2/a = 0$; and (12), $k + 1/b = 0$.

Note 2.—In the foregoing solution I have utilized the notation employed in the chapter on the Calculus of Variations in TODHUNTER'S *Integral Calculus*, fifth edition.

374. Proposed by C. N. SCHMALL, New York City.

Show that, on a *Mercator's Chart*, a great circle of a sphere of radius r_1 will be represented by a curve whose equation is of the form

$$c(e^{y/r} - e^{-(y/r)}) = 2 \sin \left(\frac{x}{r} + \theta \right).$$

I. SOLUTION BY ELIJAH SWIFT, University of Vermont.

If the latitude and longitude on the above sphere be denoted by the letters φ and θ respectively, θ varying from 0° to 360° , and φ from -90° to $+90^\circ$; then if axes be taken with origin at the center of the sphere with the xy -plane as the plane of the equator, and if longitude be measured from the x -axis, we have for any point on the sphere

$$x = r \cos \varphi \cos \theta, \quad y = r \cos \varphi \sin \theta, \quad z = r \sin \varphi.$$

The equation of a great circle is obtained by substituting these values in the equation of any diametral plane, $Ax + By + Cz = 0$, and is

$$(1) \quad A \cos \varphi \cos \theta + B \cos \varphi \sin \theta + C \sin \varphi = 0.$$

The sphere is mapped on a *Mercator's Chart* by taking a cylinder tangent to the sphere along the equator and projecting a meridian ($\theta = \text{const.}$) on a generating line of the cylinder.

Any point on the sphere on this meridian has for its image on the chart a point on the corresponding generating line at a distance $r \log \tan (\pi/4 + \varphi/2)$ from the equator.

When we develop the cylinder on a plane, we can choose axes in that plane so that the coördinates of this point are

$$x = r\theta, \quad y = r \log \tan \left(\frac{\pi}{4} + \frac{\varphi}{2} \right).$$

Solving these equations for θ and φ , $\theta = x/r$, $\varphi = 2 \arctan (e^{y/r}) - \pi/2$. Substituting these values in (1), we obtain

$$A \cos \frac{x}{r} + B \sin \frac{x}{r} + C \tan \left\{ 2 \arctan e^{y/r} - \frac{\pi}{2} \right\} = 0,$$

which reduces at once to the form given, if we let $c = -C/\sqrt{A^2 + B^2}$, and $\sin \theta = A/\sqrt{A^2 + B^2}$.